

Use Sites

Nursery, Landscape beds, Non-crop areas, Ornamental beds and Containers, Conifer Trees and other mulch beds.

Effective Against

Barnyardgrass, Bittercress, Common chickweed, Crabgrass, Dandelion, Foxtail, Goosegrass, Groundsel, Henbit, Marestail, Musk Thistle, Oxalis, Ragweed, Spurge



CRABGRASS



DANDELION







Active Ingredient

2% Trifluralin 0.5% Isoxaben

Packaging

50 lb bag

Formulation

Granular herbicide



T/I 2.5 G

Proven Performance

Up to 8-months of selective pre-emergent control of more than 100 broadleaf weeds and grasses with a single application. Performance proven on field- and container-grown ornamentals.

Mode of Action

Pre-emergent herbicide combining a cellulose biosynthesis inhibitor to control broadleaf weeds and a seedling growth inhibitor(dinitroaniline) herbicide.

Application Instructions

- Remove or incorporate all weed residue, pruning's, crop residue, or other plant trash before application
- 2. Soils surface should be smooth and free of clods when application of T/I 2.5G is made.
- If wind conditions favor drift, do not apply.Granules that drift outside of treated area may result in injury to non-target plants.
- 4. For best results, apply under mulch beds .

Spreader Settings

Spreader settings are approximate and can vary with ground speed, pattern, adjustments, wear, and changes in the manufacturing of the equipment. Therefore, you should ALWAYS calibrate your spreader before applying any product.

	Settings for Warren T-7 II Spreader ¹		
	Target Rate of Quali-Pro T/I 2.5G		
Speed (mph)	100 lbs/acre	150 lbs/acre	200 lbs/acre
2.0	2 1/2	3 1/4	3 3/4
2.5	3	3 1/2	4 1/4
3.0	3 1/4	4	5

¹ Warren spreader settings are displayed in half-number increments only. The suggested settings that indicate quarter settings can be accomplished by placing the calibration arm between the half number marks on the spreader.

