

ALL KLEER

ESSENTIAL OIL INSECTICIDE

- For Indoor & Outdoor use
- Kills ants, cockroaches, scorpions, spiders and other listed pests
- Suitable for use on edible plants, herbs, and vegetables
- Shake well

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

NET CONTENTS: **ONE GALLON**



**Control
Solutions Inc.**
ADAMA PROFESSIONAL SOLUTIONS

ALL KLEER

ESSENTIAL OIL INSECTICIDE

This product qualifies for exemption from EPA registration under Federal Insecticide, Fungicide and Rodenticide Act (FIFRA).

ACTIVE INGREDIENTS:

Clove Oil	9.0%
Geraniol	6.0%
Corrmint Oil	3.0%
Sodium Lauryl Sulfate	2.5%
OTHER INGREDIENTS*:	79.5%
TOTAL:	100.00%

*Water, Oleic acid, Polyglyceryl oleate, Isopropyl myristate

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements, Directions for Use, and Storage and Disposal. See below for additional Precautionary Statements



NET CONTENTS: 1 GALLON



Manufactured by:
Control Solutions Inc.
5903 Genoa-Red Bluff
Pasadena, Texas 77507
281-892-2500

FIRST AID

IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF INGESTED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product (including health concerns or medical emergencies), you may also contact SafetyCall® International at **1-844-685-9173**, twenty-four (24) hours per day, seven (7) days per week.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May cause eye or skin irritation. Avoid contact to eyes, skin, and clothing. May cause an allergic skin reaction. Thoroughly wash face, hands and any exposed skin after handling. Do not eat, drink, or smoke when using this product. Take off contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

While handling concentrate, applicators and other handlers must wear:

- Waterproof gloves
- Long-sleeved shirt and long pants; and
- Shoes plus socks

Follow PPE manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Surface water quality may be impacted if this product is not used according to label directions. Do not contaminate water with concentrate or rinsate.

RESTRICTIONS:

- Do not apply undiluted.
- Do not make applications to, or contaminate via drift, food preparation surfaces.
- Do not allow applications to contact people or pets.
- Phytotoxicity to treated plants may occur if this product is applied at higher than the labeled rates.
- Not for aquatic use, do not apply directly to water.
- Allow material to dry completely before people or pets are allowed to re-occupy the treatment area.

DIRECTIONS FOR USE

Read entire label prior to use and use in accordance with label instructions.

Shake well before using.

USE SITES: Includes, residential and non-residential landscapes, non-food and food areas in and around institutional, public, commercial, and industrial buildings, indoor and outdoor areas of restaurants, breweries, cafeterias, bakeries, salons, theaters, drive-in theaters, food trucks, parks, campgrounds, recreational areas, apartment buildings, day care facilities, playgrounds, nursing facilities, hospitals, schools, hotels, laboratories, lavatories, kitchens, offices, stores, storage areas, kennels, zoos (including animal enclosures), dog parks, boat houses, boat docks, marinas, mausoleums, modes of transportation (passenger vehicles, RVs, boats, buses, trucks, trailers, storage containers, railcars, marine vessels, cargo vessels), prison facilities, jail cells, lawns, turfgrass, golf courses, drains and sewers, garbage areas/rooms, ornamental plants, flowers, trees, shrubs, landscaped areas, xeriscapes, indoor plantscapes, interiorscapes, house plants, potted plants, containerized plants, mulched areas, fence lines, athletic fields/parks (including trees, shrubs, flowers, evergreens, foliage plants and groundcovers) patios & outdoor seating, and non-food residential greenhouses. Also use on surface perimeters, around foundations and walls, pest harborage and breeding areas, eaves, around windows and doors, loading docks, under carports/decks, and other manmade structures or places pests are observed.

This product may be applied to edible/consumable plants, including vegetable plants, fruit trees, herbs, tobacco, and mushroom houses, greenhouses, nurseries, shadehouses, and hydroponic facilities.

KILLS: Ants (including Fire Ants), Aphids (including Melon Aphids), Bed Bugs, Beetles (including Lady Beetles), Carpet Beetle larvae, Chinch bugs, Cockroaches (including German Cockroaches), Crickets (including House Crickets), Mealybugs, Millipedes, Mites (including Two-Spotted Spider Mites), Mosquitoes, Pillbugs, Scorpions, Silverfish, Spiders (including Wolf Spiders), Spotted Lanternflies, Springtails, Stored Product Beetles (including Cigarette Beetles), Ticks (including Lone Star Ticks), and Thrips

Mixing Instructions:

To prepare spray solution: fill a clean tank with half the required amount of water or oil, followed by the desired amount of All Kleer, and then fill the tank with the remaining amount of water or oil. Mix thoroughly.

This product may be tank mixed with other insecticides including insect growth regulators, such as Pivot 10 (EPA Registration Number 53883-280) or Tekko Pro (EPA Registration Number 53883-335). Observe and follow all restrictions, precautions, and

Directions for Use which appear on the labels of these products. This product may also be combined with adjuvants. Physical compatibility may vary with different combinations of products. Compatibility tests can be prepared on a small-scale for combinations that have not been previously tested.

Precautions:

- On sensitive surfaces such as flat painted surfaces, plastics, fabrics, vinyl and wood floors, make an application to an inconspicuous area before making more comprehensive treatments to test for damage or staining.
- Phytotoxicity may occur to treated sensitive or stressed plants at higher application rates. Before application to entire plant, test a small area, flower, or leaf first and watch for signs of sensitivity. Symptoms typically appear within 24 hours.
- Mix only enough for immediate use.

Application Sites	Amount of this product per 1,000ft ²	Application Instructions
Indoor Use: Including Food and Non-Food Areas of Food Handling Establishments	1 - 2 fluid ounces Heavy Pest Pressure: 3 fluid ounces	This product may be applied using hand-held, backpack-type, compressed air, ultra-low volume foggers, or other liquid application equipment. Combine this product with desired amount of water or oil to treat approx. 1,000 ft². It is recommended to use a minimum of 1 gallon of water or oil. Shake or agitate to thoroughly mix. To control accessible, exposed stages of listed pests, thoroughly apply where these insects are usually found. Contact as many pests as possible. Pay special attention to cracks and crevices, corners, and other harborage sites. For fogging applications apply 1-3 fluid ounces of concentrate per 1,000 cubic feet. For pest-specific rates and instructions, see Indoor General Pest Control section below. Allow material to dry before people or pets are allowed to re-occupy the treatment area. Re-apply as reinfestations warrant.
Outdoor Use: Fogging and Misting Applications	1 - 2 fluid ounces Heavy Pest Pressure: Up to 6 fluid ounces	Suitable for use with backpack mister/blowers and ULV foggers. Combine this product with desired amount of water or oil to treat approx. 1,000 ft². It is recommended to use a minimum of 1 gallon of water or oil. Shake or agitate to thoroughly mix. More than 1 gallon of water or oil might be needed, depending upon vegetation density in the treatment area. Re-apply as reinfestations warrant.

Application Sites	Amount of this product per 1,000ft ²	Application Instructions
Outdoor Use: Perimeter and Broadcast applications, Including Turf, Lawns, Mulched Areas, Xeriscape Areas, and Landscape Ornamentals	1 - 2 fluid ounces Heavy Pest Pressure: Up to 6 fluid ounces	This product may be applied using hand-held, hose-end, backpack-type, compressed air, mister/blowers, high volume, or other sprayers. Combine this product with desired amount of water or oil to treat approx. 1,000 ft². It is recommended to use a minimum of 1 gallon of water or oil. More than 1 gallon of water or oil might be needed, depending upon vegetation density in the treatment area. Shake or agitate to thoroughly mix. For outdoor perimeter applications, apply in a band up to 15 feet adjacent to and around the structure where pests may be active or may enter. For broadcast lawn applications, treat desired areas with approximately 1 gallon of spray solution per 1,000 ft². For applications to landscape ornamentals, use enough spray solution to completely penetrate the leaf canopy and cover all leaves until wet without excessive runoff. Use caution when applying to open blooms. Spraying should begin when insects first appear. Do not wait until plants are heavily infested. Repeat as needed but not more than once daily. Allow material to dry before people or pets are allowed to re-occupy the treatment area. Re-apply as reinfestations warrant.
Outdoor Use: On and Around Boat Docks	1 - 2 fluid ounces Heavy Pest Pressure: Up to 6 fluid ounces	To control listed pests, dilute 1-2 fluid ounces of concentrate per 1 gallon of water or oil and make applications to the dock area. In situations with heavy pest pressure, the rate may be increased up to 6 fluid ounces per gallon. Do not treat directly into water. Re-apply as reinfestations warrant.
Livestock and Companion Animal Facilities (barns, stables, poultry houses, swine buildings, livestock sheds, runs, and other farm buildings)	1 - 2 fluid ounces Heavy Pest Pressure: Up to 6 fluid ounces	This product may be applied using backpack mister, fogger, hand-held, hose-end or backpack-type compressed air sprayers. Combine this product with desired amount of water or oil to treat approx. 1,000 ft². It is recommended to use a minimum of 1 gallon of water or oil. Shake to thoroughly mix. Remove animals from the treatment area prior to application. Make applications to surface perimeters, around foundations,

Application Sites	Amount of this product per 1,000ft ²	Application Instructions
		inside enclosures, and areas where pests congregate, rest, or breed. Allow material to dry before animals are allowed to re- occupy the treatment area. Re-apply as reinfestations warrant.
Foliar Application to Indoor and Outdoor Grown Ornamental Plants, Edible Plants, and Crops, including those in Greenhouses, Nurseries, and Hydroponic Facilities	1 - 2 fluid ounces per 1,000 square feet (43-86 fluid ounces per acre) Heavy Pest Pressure: 4 fluid ounces per 1,000 square feet (172 fluid ounces per acre)	Apply by appropriate equipment in sufficient water for thorough coverage to foliage. It is recommended to use approximately 2 gallons of spray solution per 1000 ft² or 100 gallons per acre. Mix only enough for immediate use. Use on ornamental plants including roses, flowers, shrubs, small trees, and on garden fruits, herbs, and vegetables, including tobacco, melons, berries, and nut trees. Use enough spray solution to completely penetrate the leaf canopy and cover all leaves until wet without excessive runoff. Use caution when applying to open blooms. Repeat applications as needed, 5-to-7-day intervals recommended. Spraying should begin when insects first appear. Do not wait until plants are heavily infested. Repeat as needed but not more than once daily. Hard to control pests may require additional applications. Exposure to intense light, heat, or humidity following application may increase the potential for phytotoxicity. Minimize heavy applications by applying in such a manner that allows the spray to dry within 2 hours. Applications are best tolerated in the early morning or evening when temperatures are cooler. Some plants may be sensitive to essential oils under certain conditions. Avoid use on newly planted cuttings or plants stressed by drought. Avoid applications to plants in direct sunlight when temperatures exceed 90°F. Avoid application to sensitive blooms. Allow material to dry before people are allowed to re-occupy the treatment area. Once material is dry, plants may be harvested.

Application Sites	Amount of this product per 1,000ft ²	Application Instructions
Dip Application to Indoor and Outdoor Ornamental, Edible Plants and Crops in Greenhouses, Nurseries and Hydroponic Facilities	1.5 fluid ounces	Mix 1.5 fl. oz. per gallon of water. This product may be used to control pests when applied as a dip of unrooted cuttings or foliage (above-ground portions) of rooted cuttings. Clean and disinfect the dipping tank and equipment before a new dip solution is prepared. Prepare only as much dip solution as can be used in one day. If plant pathogens are a concern, prepare a new dip solution regularly throughout the day. Ensure full coverage by dipping for approximately 10-30 seconds while moving the plants around in the solution. Agitate dip solution throughout use. Certain plants may show phytotoxicity symptoms post-dipping. Prior to treating entire crop, test a small portion of the crop for sensitivity. Do not dip stressed/wilted cuttings or transplants.
Grazed Pastures, Hay Fields, Rangeland, Grass grown for Silage or Hay, Grass Fodder, Grass Forage	1 - 2 fluid ounces (43-86 fluid ounces per acre) Heavy Pest Pressure: 4 fluid ounces (172 fluid ounces per acre)	Ground application: Apply when insects appear in sufficient spray volume to obtain full coverage of the foliage or target area. For pests in thatch more volume may be needed. Air application: Apply in a minimum of 2 gallons per acre of finished spray or sufficient volume to obtain full coverage of the target area. Use higher rates or increased pest pressure. Monitor insect populations to determine timing and frequency of applications. Pasture and rangeland grass: May be used for grazing or cut for forage 0 days after application. Do not cut grass to be dried and harvested for hay until the application has dried.

Indoor General Pest Control

This product may be applied to the interior of residential, commercial, and industrial facilities in both food and non-food areas. Before application, ensure food is covered or removed. Apply at 1-2 oz dilution rate. Use in locations where pests are present. Apply as a cracks and crevice, voids or broadcast applications to floors and carpets where pests may live or hide. Common harborage areas include around water or utility pipes, baseboards and floorboards, carpeted floors, cabinets and drawers, behind dishwashers/refrigerators/appliances, door and window frames, voids, and other pest harborage areas. Contact as many pests as possible. Do not apply to electrical voids. Do not apply undiluted.

Ants: Apply product at the 2 fl oz per gallon of water or greater dilution rate, depending on pest pressure. Make applications directly to mounds, trails, and cracks where ants are traveling as a spot and/or crack and crevice treatment.

Bed Bugs: Apply product at the 2 fl oz per gallon of water or greater dilution rate, depending on pest pressure. Make applications to cracks and crevices where bed bugs may live. Common bed bug harborages include behind wall-mounted headboards, picture frames and floorboards, screw holes and other crevices in the bed frame, and rugs/carpet. Treat the folds, tufts, and seams of mattresses and upholstered furniture. Often harborages are beyond the bed, such as near couches and sofas. Carpets and rugs may also be treated. Avoid over saturating any surface.

Stored Product Pests: For cigarette beetles and other stored product pests, remove food and food sources before treating pantries, cabinets, drawers, and other stored product pest harborages. Do not treat food directly. Apply product at the 2 fl oz per gallon of water or greater dilution rate, depending on pest pressure.

Spiders: Apply product at 2 – 6 fl oz per gal of water.

Scorpions: Mix 3 - 6 fl oz per gallon of water and apply to areas where pests are observed. Treat baseboards, flooring, hardwood floors, carpeted floors, rugs, pet resting areas, window and door frames, attics and crawlspaces.

Ticks: Use 1 – 6 fl oz per gallon of water and apply to areas where ticks are observed or suspected. Wash pet bedding with hot water. Treat crawlspaces, baseboards, flooring, hardwood floors, carpeted floors, rugs, pet resting areas, window and door frames, draperies, and furniture folds, tufts and seams. Avoid over saturating any surface.

Food Handling/Processing Establishments

When using as a fog, apply 1-2 fl oz of concentrate per 1,000 cubic feet. Ensure food and food products are removed and cover all food processing surfaces and equipment. Those pieces of equipment that cannot be removed or covered, clean and rinse after application before operation begins. Do not apply as a fog when the facility is in operation; all employees and non-employees must exit the application area. For oil dilutions in food handling establishments, ensure the deodorized petroleum oil conforms to CFR 21, section 172.884. Use caution as fogging oils may create slippery conditions. Ventilate treatment area before re-occupying.

When used as a spot or crack and crevice treatment, the facility may remain in operation.

Outdoor Premises Fogging and Misting Applications:

For best results, treat ornamental plants, hedges, shrubs, and trees where pests may congregate, rest, or breed, including under the leaves, where adult mosquitoes and ticks or other listed pests hide during the day. Treat under shrubs, wood piles, leafy areas, vining ground cover, landscaped/mulched areas, hedges, and anywhere that is shaded and moist. For tick control, apply around edges of lawns, under porches, where siding meets foundations, cracks, fences, and areas where pets or wildlife may travel or rest. Vegetation around stagnant pools, marshy areas, and shorelines may be treated. Do not spray directly on ponds of any kind, lakes, rivers, or other bodies of water. Make applications to covered porches, carports, window frames, doors, garage doors, patios, carports, shutters, edges of siding, open crawlspaces, waste receptacles, dumpsters, inside irrigation control boxes, utility boxes, under decks, and eaves. Broadcast applications to turf or lawns may be made.

Outdoor Perimeter and Broadcast Use:

Apply around decks, garage areas, sill plates, storage sheds, trunks of woody plants, mulch, fence lines, xeriscaped/landscaped/mulched areas, lawns, turf, artificial turf, surface perimeters, and other areas where pests may congregate. Apply to window frames, doors, garage doors, patios, covered walkways, eaves, carports, around foundations and walls, and other areas where pests congregate. Contact as many pests as possible.

Scorpions: Apply 3 - 6 fluid ounces per 1,000 ft² as a coarse, low pressure application where scorpions are known to live, breed, hide, or enter the structure. For best results, apply higher volumes of water to penetrate to harborage. It is recommended to remove potential harborage including rock piles, lumber, firewood and other areas where scorpions are known to rest. To improve longevity of control, a tank mix with an EPA registered residual product is recommended.

Spiders: Apply 2 - 6 fluid ounces as a course, wet spray to areas where pests are observed. Treat around doors & windows, attics, crawl spaces, eaves, corners, cracks & crevices, openings, entry points, and harborage areas. Contact as many pests as possible. Repeat as needed.

Ticks: Apply 1 - 6 fluid ounces per 1,000 ft² as a coarse, low pressure application where ticks are known or suspected. Make applications to shaded areas, under trees, areas where animals rest, under decks, retaining walls, and other pest harborage areas. This product may be tank mixed with other insecticides, including insect growth regulators, or adjuvants to enhance control.

Outdoor Automatic Misting Systems:

To control adult mosquitoes and other listed pests, add water and concentrate to the tank to achieve desired dilution rate. The recommended rate is 1-2 fluid ounces per 1,000 ft² of treated area with a maximum of 6 fluid ounces per 1,000 ft². Agitate mixture prior to misting. Dilutions may be stored for extended periods of time, not to exceed 1 month (30 days). Follow the system manufacturer's directions for maintenance and settings. Wait for the mist to dissipate before people or pets enter the treated area.

Site	Section / Listed Pests*	Standard Application Rate (Fluid Ounces of Product per Gallon)**	Dilution Application
Indoor	Bed Bugs	2	Minimum 1 gal per 1,000ft ²
Indoor	Ants	1 - 2	Minimum 1 gal per 1,000ft ²
Indoor	Stored Product Pests	2	Minimum 1 gal per 1,000ft ²
Indoor	Spiders	2 - 3	Minimum 1 gal per 1,000ft ²
Indoor	Scorpions	3 - 6	Minimum 1 gal per 1,000ft ²
Indoor	Food Handling Establishments	1 - 3	Minimum 1 gal per 1,000ft ²
Outdoor	Fogging & Misting	1 - 2	Minimum 1 gal per 1,000ft ²
Outdoor	Ticks	1 - 2	Minimum 1 gal per 1,000ft ²
Outdoor	Scorpions	3	Minimum 1 gal per 1,000ft ²
Outdoor	Spiders	2	Minimum 1 gal per 1,000ft ²
Outdoor	Boat docks	1 - 2	Minimum 1 gal per 1,000ft ²
Outdoor	Automatic Misting Systems	1 - 2	1 – 6 ounces of product per 1,000 ft ² of treated area
Outdoor	Perimeter, Broadcast, & Turf	1 - 2	Minimum 1 gal per 1,000ft ²
Outdoor	Grazed Pastures, Hay Fields, Rangeland	1 - 2	Minimum 2 gal per acre
Indoor & Outdoor	Livestock and Companion Animal Facilities	1 - 2	Minimum 1 gal per 1,000ft ²
Indoor & Outdoor	Ornamental and Edible plants (aphids, mites, mealybugs, thrips, etc.)	1 - 2	Minimum 2 gal per 1,000ft ² or 100 gallons per acre

*Refer to the tables and sections above for full pest- and site-specific application directions.

**For heavy infestations indoors, use up to 3 fluid ounces of All Kleer. For heavy infestations outdoors (including perimeter and broadcast applications to turf, livestock and companion animal facilities), use up to 6 fluid ounces of All Kleer. For Foliar Application to Indoor and Outdoor Grown Ornamental Plants, Edible Plants, and Crops (including those in Greenhouses, Nurseries, and Hydroponic Facilities) and to grazed pastures, 4 fluid ounces per gallon for heavy infestations may be used.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. **DO NOT** store this product above 100°F for extended periods of time. Avoid extremes in temperatures. If container is damaged or if a spill has occurred, use product immediately, or dispose of product as indicated below.

PESTICIDE DISPOSAL: **DO NOT** pour or dispose of down the drain or sewer. **CONTAINER HANDLING:** **DO NOT** reuse or refill this container. Empty container by using the product according to the label directions. Triple rinse container (or equivalent) promptly after emptying. Offer for recycling if available or place in trash receptacle.

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

WARRANTY STATEMENT

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. **CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, **IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT** in no event shall Control Solutions or Seller be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

To the extent consistent with applicable law, Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

