

Feb 20, 2026

**QUALI-PRO**

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

**CONTRADO<sup>®</sup> SC**

Control Solutions, Inc.  
5903 Genoa Red Bluff  
Pasadena, TX 77507

Classified for  
“RESTRICTED USE”  
in New York State  
under 6NYCRR Part 326

**24(c) Special Local Need Label No. 260002**

**FOR USE IN NASSAU, SUFFOLK, KINGS, OR  
QUEENS COUNTIES IN NEW YORK STATE  
ON GOLF COURSES AND SOD FARMS ONLY**

EPA SLN No:

EPA Reg. No: 53883-534

**FOR DISTRIBUTION AND USE ONLY IN  
NEW YORK**

## For foliar and systemic control of insect pests in or on golf courses and sod farms.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This state-specific section 24(c) labeling must be in possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Contrado Insecticide (EPA Reg. No. 53883-534) and this label.

### USE RESTRICTIONS

- **DO NOT** let people and pets enter the treated area until treatment has dried.
- **DO NOT** use for commercial grass seed production.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product with aerial application equipment.
- **DO NOT** apply more than 20 fl. oz. Contrado<sup>®</sup> SC insecticide (equivalent to 0.26 lb of any chlorantraniliprole-containing products) per acre per calendar year to plants produced or grown outdoors.
- **DO NOT** apply more than a single application per year.

The following restrictions are required to permit use of this product in the State of New York:

- **Golf course greens and tee boxes:**
  - DO NOT apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- **Golf course fairways:**
  - DO NOT apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- **Other use sites:**
  - DO NOT apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

| APPLICATION RATES  |                                |                          |                       |  |
|--|--------------------------------|--------------------------|-----------------------|--|
| Target Pest  | fl. oz. product/<br>1000 sq ft | fl. oz. product/<br>Acre | Lbs. A.I.<br>per acre | Application Instructions   |
| Armyworms<br>Cutworms<br>Sod Webworms  | 0.05 – 0.09                    | 2 – 4                    | 0.026 –<br>0.052      | Apply preventatively or after caterpillars have been observed.<br>To ensure best control, delay water (irrigation) or mowing for 24 hours after application.<br>If the area being treated is maintained at a mowing height greater than one inch, the higher rate listed may be required during periods of high pest pressure.                 |
| Billbugs   | 0.18 – 0.46                    | 8 – 20                   | 0.104 –<br>0.26       | Apply when overwintered adults are first observed, This will usually occur in late April or early May for cool-season turfgrass. The higher rates listed may be required for applications made after early May or when billbug species other than bluegrass billbug are present.<br>Applications at this time also provide white grub control. |
| European crane fly larvae  | 0.18 – 0.37                    | 8 – 16                   | 0.104-<br>0.208       | Apply between September and November. The higher rate listed may be required to achieve control when applications are made in November.  |
| White grub larvae, including: <ul style="list-style-type: none"> <li>• <i>Aphodius spp.</i></li> <li>• Asiatic garden beetle</li> <li>• Black turfgrass ataenius</li> <li>• European chafer</li> <li>• Green June beetle</li> <li>• Japanese beetle</li> </ul> | 0.18 – 0.37                    | 8 – 16                   | 0.104 –<br>0.208      | Apply from early April to early September for preventative control. The higher rate listed may be required for early-season (April and May) applications where a long residual is required or in late August or early September when less-sensitive mid-instar grubs   |

|  |  |  |  |
|--|--|--|--|
| <ul style="list-style-type: none"> <li>• May/June beetles (<i>Phyllophaga spp.</i>)</li> <li>• Northern masked chafer</li> <li>• Oriental beetle</li> <li>• Southern masked chafer</li> </ul>  |  |  | <p>are present at the time of application. Irrigate turf within 48 hours after application or allow rainfall to move the product into the soil. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping, or other methods.</p> |
| <p><b>USE RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>• Maximum Single Application Rate <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 20 fl oz of product per acre (0.26 lb ai per acre).</li> </ul> </li> <li>• Maximum Annual Application Rate <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 20 fl oz of product per acre (0.26 lb ai per acre).</li> </ul> </li> <li>• <b>DO NOT</b> enter or allow others to enter treated areas until sprays have dried.</li> </ul> |  |  |  |